

**LTBI: New Options for Short Course Treatment References**

1. Comstock, G. W., and S. H. Ferebee. **1970**. How much isoniazid is needed for prophylaxis? *Am. Rev. Respir. Dis.* 101:780–782.
2. Comstock, G. W. **1999**. How much isoniazid is needed for prevention of tuberculosis among immunocompetent adults? *Int. J. Tuberc. Lung Dis.* 3:847–850.
3. Ferebee, S. H. **1970**. Controlled chemoprophylaxis trials in tuberculosis: a general review. *Adv. Tuberc. Res.* 17:28–106.
4. Hong Kong Chest Service, Tuberculosis Research Centre, Madras, and British Medical Research Council. **1992**. A double-blind placebo controlled clinical trial of three antituberculosis chemoprophylaxis regimens in patients with silicosis in Hong Kong. *Am. Rev. Respir. Dis.* 145:36–41.
5. Ormerod, L. P. **1998**. Rifampicin and isoniazid prophylactic chemotherapy for tuberculosis. *Arch. Dis. Child* 78:169–171.
6. Joint Tuberculosis Committee of the British Thoracic Society. **1998**. Chemotherapy and management of tuberculosis in the United Kingdom. *Thorax* 53:536–548.
7. CDC. **2000**. Targeted tuberculin testing and treatment of latent tuberculosis infection. *MMWR*
8. Martinson, G, Barnes G, et al. **2011**. New Regimens to Prevent Tuberculosis in Adults with HIV Infection (Soweto study): *N Engl J Med*;365:11-20
9. Samandari T, Agizew T, etc al. **2011**. 6-month versus 36-month isoniazid preventive treatment for tuberculosis in adults with HIV infection in Botswana: a randomised, double-blind, placebo-controlled trial. *Lancet*; 377:1548-1550.